

© 2007 EcoHealth Journal Consortium

#### Review

# Is Marine Mammal Health Deteriorating? Trends in the Global Reporting of Marine Mammal Disease

Frances M. D. Gulland<sup>1</sup> and Ailsa J. Hall<sup>2</sup>

Abstract: A recent rise in the reporting of diseases in marine organisms has raised concerns that ocean health is deteriorating. The goal of this study was to determine whether or not there has been a recent deterioration in marine mammal health by investigating the trends in disease reports over the past 40 years (categorized by the method of study, the species affected, and the etiology of the disease) and by exploring the changes in frequency of mass mortality events among marine mammals reported in the United States since 1978. The number of papers on marine mammal disease published each year has increased since 1966, although the annual publication rate appears to have stabilized since  $\sim$ 1992. Those published in the 1960s and 1970s were largely about helminth and bacterial disease, those investigating viruses emerged in the late 1970s and increased in the 1980s and 1990s, whereas protozoal diseases and harmful algal toxins were largely not reported until the 1990s. The annual number of mass mortality events in the U.S. approximately doubled between 1980 and 1990 but since 2000 has been between seven and eight events per year. Causes of mass mortality events have included biotoxins, viruses, bacteria, parasites, human interactions, oil spills, and changes in oceanographic conditions. Events due to biotoxins appear to be increasing, but the change in the frequency of mass mortality events from other causes over the past 40 years cannot be determined from the available published literature due to changes in marine mammal abundance, inconsistencies in effort and extent of resources for pathological investigation, and advances in technology that have allowed improved detection of pathogens and toxins in more recent years. To ensure future information on the true incidence of marine diseases and their underlying causes is more reliable, specific and directed marine health monitoring programs, well-equipped stranding networks, and dedicated diagnostic laboratories are needed.

Keywords: disease, epidemiology, marine mammal, mortality, morbidity

#### Introduction

A recent rise in the reporting of diseases in marine organisms has raised concerns among scientists, politicians,

Published online: April 12, 2007

Correspondence to: Frances M. D. Gulland, e-mail: gullandf@tmmc.org

managers, and the public that ocean health is deteriorating (Epstein, 1996; Geraci et al., 1999; Harvell et al., 1999; Gardner et al., 2003). Marine diseases, particularly those that cause large-scale die-offs, can alter population distribution and abundance, and cause major regime changes within marine communities (Harvell et al., 2004; Kim et al., 2005). Furthermore, threats to ocean health can directly

<sup>&</sup>lt;sup>1</sup>Veterinary Science Department, Center for Marine Animal Health and The Marine Mammal Center, 1065 Fort Cronkhite, Sausalito, CA, 94965, USA <sup>2</sup>Center for Marine Animal Health and Sea Mammal Research Unit, University of St. Andrews, St. Andrews, Fife, UK

and indirectly impact human health, and many emerging diseases of wildlife are zoonotic (National Research Council, 1999; Knap et al., 2002; Knowlton, 2004; Cunningham, 2005). However, whether the increase in reports represents a real and widespread degeneration in the health of marine life is unclear (Harvell et al., 1999; Lafferty et al., 2004; Ward and Lafferty, 2004). The lack of information on the true incidence of marine mammal diseases (i.e., the rate at which new cases of the disease are occurring during a specified period) and their underlying causes are largely due to the lack of specific and directed marine health monitoring (Harvell et al., 2002; Kim et al., 2005). This is probably a consequence of the historical focus on domestic animal health rather than wildlife disease, and the emphasis on farm animal economy rather than conservation.

Marine mammals are particularly difficult marine organisms to study due to their aquatic, often pelagic, and non-sedentary life styles, long generation times, large body size and protected status. However, when marine mammals do wash ashore (strand) sick or dying, they command considerable attention and concern. There is, thus, an interest from the public, as well as from scientists and managers, to determine the state of marine mammal health and to identify the anthropogenic influences (such as exposure to pollutants and terrestrial pathogens, increased ocean noise and boat harassment) on the well-being of these animals. The stranding of beaked whales in association with navy exercises has raised awareness of direct effects of human activities on marine mammal health (Fernández et al., 2004; Cox et al., 2006). Indirect effects are less obvious, but there is no doubt that overt morbidity and mortality among marine mammals have resulted from terrestrial pathogens spreading to the ocean, as well as from harmful algal blooms and epidemics of virulent viruses and bacteria (Geraci et al., 1999; Miller et al., 2002). Although the role of anthropogenic factors and climate change in these die-offs is uncertain, there is a compelling body of evidence indicating that persistent organic pollutants can affect marine mammal immunity (DeSwart et al., 1994; Ross et al., 1995; Hammond et al., 2005) and the exposure of individuals to these compounds may have played a role in some viral mass mortality events (Hall et al., 1992; Aguilar and Borrell, 1994). Furthermore pathogens of terrestrial origin, such as Toxoplasma gondii and antibioticresistant bacteria are causing disease in marine mammals, and appear to originate in fresh water run-off (Miller et al., 2002; Stoddard et al., 2005). Thus anthropogenic effects on the ocean may be adversely impacting marine mammal


health, causing an increase in disease outbreaks and deterioration in health.

Direct data on conventional indices used to quantify health in humans and domestic animals (hematology, serum biochemistry, and immune function markers) are relatively sparse for marine mammals, and there are insufficient published data over several decades to assess changes in health using these reports. Instead, we use frequency of disease outbreaks as indicators of health status. Data were used from two sources. Firstly, we investigate the trends in disease reports published in the scientific literature over the past 40 years, categorized by the method of study, the species affected, and the type of organism or toxin involved; secondly, we explore the changes in frequency of mass mortality events among marine mammals reported both in the United States (U.S.) since 1978, and worldwide. Data on mass marine mammal mortality events in the U.S. are considered separately not because this country is representative of the global condition, but it has a very long coastline, a wide variety of well-studied Atlantic and Pacific marine mammal species, and, under the auspices of the Marine Mammal Protection Act, has a formal and structured response to mass marine mammal mortality events, which means that effort at reporting these larger events has been relatively consistent since 1978.

#### **M**ETHODS

# Papers on Disease Published in Peer Reviewed Journals

To investigate and assess the trends in marine mammal disease, we reviewed 772 papers published in peer reviewed journals since 1966. Databases searched included PubMed, Medline, CAB abstracts, Google scholar, and ISI Web of Knowledge; search terms included marine mammal, pinniped, cetacean, disease, health, virus, bacteria, toxin, parasite, epidemic. A list of the papers (with their abstracts) included can be found at http://www.smru.st-and.ac.uk. Papers were categorized by both authors according to disease agent investigated, method used to investigate disease in marine mammals, and by species affected. Papers described diseases in individual marine mammals, as well as mass mortality events. Papers were published 1-5 years after a disease was identified, so these data reflect trends in disease reporting over time and not the absolute changes in the true dates that each disease was identified.



**Figure 1.** Number of papers on disease in marine mammals published in peer reviewed journals, 1960–2006. Note: 2006 includes papers to July only.

#### **Mass Mortality Events**


Reports of mass mortality events worldwide were obtained from the published literature, and reports of unusual mortality events in the United States were also obtained from the administrative records of the National Marine Fisheries Service (NMFS) Marine Mammal Health and Stranding Program (Silver Spring, Maryland, U.S.). These records result from stranding network members detecting sick or dead marine mammals on the beach and submitting data on each animal (including date, species, length, age class, sex, and location) to the regional stranding coordinators of the NMFS. In the U.S., large mortality events involving marine mammals are referred to in two different ways. A stranding event, in which animals usually beach alive, involving more than two marine mammals (other than mother-calf pairs) are defined as "mass strandings" (Geraci et al., 1999). These mass strandings are most common on the eastern coastline, typically involve odontocetes, and the cause is generally not identified (Geraci et al., 1999). Other mass mortalities are referred to as "unusual" mortality events as unusually large numbers of animals die. Furthermore, in an effort to improve response and investigation of these unusual mortality events, a formal federal designation of "Unusual Mortality Event" (UME), also exists, the definition being applied to an unusual event following the vote of a federally appointed group of experts that use seven criteria upon which to base their decisions. These criteria include: (1) a marked increase in the magnitude or a marked change in the nature of morbidity, mortality, or strandings when compared with prior records; (2) a temporal change in morbidity, mortality, or strandings is occurring; (3) a spatial change in morbidity, mortality, or strandings is occurring; (4) the species, age, or sex composition of the affected animals is different than that of animals that are normally affected; (5)


affected animals exhibit similar or unusual pathologic findings, behavior patterns, clinical signs, or general physical condition (e.g., blubber thickness); (6) potentially significant morbidity, mortality, or stranding is observed in species, stocks, or populations that are particularly vulnerable (e.g., listed as depleted, threatened, or endangered or declining); (7) morbidity is observed concurrent with or as part of an unexplained continual decline of a marine mammal population, stock, or species. Since 1978, the administrative record of unusual mortality events has been relatively consistent, although the formal designation of the term "unusual" has varied. Thus, for this article, we examine all records of unusual mortality events, to minimize the effect of changes in nomenclature on our ability to detect changes in frequency of disease outbreaks. Mass strandings are not considered here, as these typically have not been determined to result from disease outbreaks (Geraci et al., 1999).

#### RESULTS

# Papers on Disease Published in Peer Reviewed Journals

The number of papers on marine mammal disease published in the scientific literature each year has increased (Fig. 1), although the annual publication rate appears to have stabilized since  $\sim$ 1992. In an attempt to determine any trends in the etiologies of marine mammal disease, we categorized the published studies by the agents for the disease being investigated (viral, bacterial, helminth, fungal, protozoal, and harmful algal toxins). The trend in the different disease agents studied (as a proportion of the total number reviewed, by 3-year categories) is shown in Figure 2a and b. Those papers published in the 1960s and





**Figure 2.** Proportion of the total number of papers by 3-year categories, relating to (a) viral, bacterial, or helminth disease; (b) fungal, protozoal disease, or harmful algal blooms.

1970s were largely about helminths and bacterial disease; those investigating viruses emerged in the late 1970s and increased in the 1980s and 1990s. Fungal disease, particularly lobomycosis in dolphins, seems to have been particularly studied in the late 1970s and early 1980s, whereas protozoal diseases and the effects of harmful algal toxins on marine mammals were largely not reported until the 1990s (Fig. 2b). Although the studies must be interpreted in the light of the type of investigation and may, to some extent, be biased by any delay in publication, they appear to reflect our increased ability to detect and identify more fastidious pathogens, such as small bacteria, protozoa, and algal toxins in more recent years. As novel laboratory techniques have been introduced, the number of publications on the agents that the techniques can detect increased. For example, the introduction of polymerase chain reaction (PCR) as a routine diagnostic tool for detecting viruses in the late 1980s was associated with an increase in reports of virus infections in marine mammals, and use of high performance liquid chromatography for the detection of biotoxins was not common until the 1990s.


The scientific approaches used to study the occurrence of disease have also changed over time. Most studies are investigations on individuals and 12% of the publications were case studies, reporting disease on one animal. Few

studies have included sufficient individuals to be able to make inferences about impacts of disease at the population level, with the exception of the effect of mass mortalities. Figure 3a and b shows how the proportion of each sampling method reported (of the total number of papers) has changed since 1966. The number of studies using harvested and captive animals has steadily declined (Fig 3a). Studies where only animals by-caught in fishing nets were included are not seen until the late 1990s, but studies on stranded animals will include by-caught animals as well as those found washed ashore. Studies on stranded animals have increased, and constitute approximately 57% of the papers reviewed (Fig. 3b). Studies from live stranded and rehabilitated marine mammals have also increased, with few studies on these animals prior to 1985. Papers on mass mortalities, compared with studies on individually stranded animals, peaked in the early 1990s. This peak reflects reports from the high profile morbillivirus outbreaks in the U.S. and Europe in the late 1980s. A second peak occurs in the 2005-present year group, largely as a result of publications on the 2002 European phocine distemper virus (PDV) outbreak.

The vast majority of published studies have been carried out on the harbor seal (*Phoca vitulina*), with the bottlenose dolphin (*Tursiops truncatus*) the most commonly studied cetacean. This probably reflects the relative abundance and near-shore distribution of these two species and the fact that bottlenose dolphins have often been kept in captive display facilities where case studies (particularly on lobomycosis) have been carried out. Harbor seals have also been subject to mass mortalities in recent years and harbor porpoises (*Phocoena phocoena*), which also feature high on the list, are often stranded or by-caught. Interestingly, the distribution is about equal between seals and cetaceans, but it is clear that almost nothing has been reported about diseases in some species.

### Temporal and Spatial Trends in Unusual Mortality Events


The total number of unusual mortality events reported over time in the U.S. is shown in Figure 4a. The annual number of events approximately doubled between 1980 and 1990 but, since 2000, it has remained at between seven and eight events per year. Causes of mortality events have included biotoxins, viruses, bacteria, parasites, human interactions, oil spills, and changes in oceanographic conditions





**Figure 3.** Proportion of the total number of papers by 3-year categories, using (a) by-caught, harvested, captive, or wild captured; (b) individual strandings and mass mortalities; (c) rehabilitated marine mammals to study the occurrence of disease.

(Fig. 4b). Prior to 1996, when a manatee die-off was caused by brevetoxin, only five mass mortality events were associated with exposure to biotoxins from harmful algal blooms. Since then, 12 of 31 events in the U.S. have been associated with exposure to biotoxins (Fig 4c), and events caused by domoic acid on the West coast of the U.S. and brevetoxin on the East coast are increasing in frequency. This reflects a true increase in the frequency of harmful algal blooms around the U.S., especially over the last decade (Van Dolah, 2000).

Frequency of unusual mortality events from other causes has not obviously changed in recent years. Interestingly, influenza virus was the cause of two mortality







**Figure 4.** Marine mammal unusual mortality events in the U.S. (data from National Marine Fisheries Service database as listed in Table 1) by 2-year categories in (a) total number of marine mammal mortality events in the U.S. by year; (b) by identified causes, 1978–2005 (HAB = harmful algal blooms; other = human interaction, oil spills, and changes in oceanographic conditions); (c) in different taxa, 1978–2005.

events in 1979 and 1982, but in more recent years, influenza epidemics in seals have not been detected and morbillivirus epidemics have been the more common causes of virus-associated mass mortality events.

Some other types of mortality events, such as those along the California coast associated with El Niño and leptospirosis, also occur at regular intervals. Mortality events in the U.S. have been most frequent in Florida and California, and have been most frequently reported in bottlenose dolphins, California sea lions (*Zalophus californianus*), and manatees (*Trichechus manatus latirostris*).

Table 1.	Summary of Marine Mammal Unusual Mass Mort	ortality Events Reported 1978–2005 <sup>a</sup>		
Year	Species and no. affected	Location	Cause and comments	References
1978	Hawaiian monk seals Monachus schauinslandi (50)	Northwest Hawaiian Islands, U.S.	Ciguatoxin and maitotoxin suspected	Gilmartin et al., 1980
1979–1980	Ĭ	Cape Cod, Massachusetts, U.S.	Influenza A (a mycoplasma was concurrently isolated from these seals)	Geraci et al., 1982
1982 1982	Harbor seals <i>Phoca vitulina</i> Florida manatees <i>Trichechus</i>	Cape Cod, Massachusetts, U.S. Southwest Florida, U.S.	Influenza A Brevetoxin	Hinshaw et al., 1984 O'Shea et al., 1991
1983	Several pinniped species (especially California and Galapagos sea lions Zalophus californianus); Galapagos	West coast of U.S.; Galapagos	El Niño	Trillmich and Ono, 1991
1984	tut seats (motasmus) California sea lions Zalophus	California, U.S.	Leptospirosis	Dierauf et al., 1985
1987 1987	Sea otters Enhydra lutris (34) Humpback whales Megaptera	Kodiak Island, Alaska, U.S. Massachusetts, U.S.	Saxitoxin Saxitoxin	DeGange and Vacca, 1989 Geraci et al., 1989
1987–1988	ВС	New Jersey, Delaware, Virginia, North Carolina, South Carolina, Georgia, Florida, U.S.	Morbillivirus. Brevetoxin was detected in dolphins, its role in the event is unclear	Scott et al., 1988; Geraci, 1989; Lipscomb et al., 1994; Duignan et al., 1996; Schulman et al., 1997; Eriedlaander, 2000: McTellan et al., 2002
1987–1988 1988 1989 1990	Baikal seals <i>Phoca siberica</i> (80–100,000)  Harbor seals <i>Phoca vitulina</i> (~18,000)  Sea otters <i>Enhydra lutris</i> (3500–5000)  Striped dolphins <i>Stenella coeruleoalba</i> (550)  Bottlenose dolphins  Tursiops truncatus (146)	Lake Baikal Northern Europe; U.K. Alaska, U.S. Mediterranean Sea Texas, Louisiana, Mississippi, Alabama, U.S.	Canine distemper virus (CDV) Phocine distemper virus (PDV) Exxon Valdez oil spill Dolphin morbillivirus (DMV) Unknown. Unusual skin lesions observed	Grachev et al., 1989; Osterhaus et al., 1989 Kennedy et al., 1988; Osterhaus et al., 1988 Loughlin, 1994 Aguilar and Raga, 1993; Domingo et al., 1992 B. Brown memo to N. Foster, March 2, 1990; Kuehl and Haebler, 1995; Hansen, 1992; Medway report to Fox, June 29, 1990

Table 1.	. Continued			
Year	Species and no. affected	Location	Cause and comments	References
1991	Harbor seals <i>Phoca vitulina</i> (34)	New Jersey, U.S.	Unknown. Erysipelothrix rhusiopathiae cultured from 4 seals, poxvirus present in 1, saxitoxin negative	Informational memo J. A. Knauss, May 10, 1991
1991	California sea lions <i>Zalophus</i> californianus (160)	California, U.S.	Leptospirosis	Gulland et al., 1996a
1991	Bottlenose dolphins Tursiops truncatus (30)	(Sarasota) Florida, U.S.	Unknown	Working Group files, NMFS
1992	Harbor seals <i>Phoca vitulina</i> (29); Harbor porpoises <i>Phocoena phocoena</i> (5); Rough toothed dolphin <i>Steno bredanensis</i> (1)	Oregon, Washington, U.S.	Unknown	H. Braham memos to W. Aron Oct. 9, 1992; Oct. 16, 1992; and Dec. 1992
1992	Dugong Dugong dugong (100)	Queensland, Australia	Starvation	Preen and Marsh, 1995
1992	Common dolphins Delphinus delphis (118)	Southwest England, U.K.	Fisheries interaction	Kuiken et al., 1994
1992	Phocid seals (24)	Maine, Massachusetts, Connecticut, U.S.	Unknown. Morbillivirus	Geraci et al., 1993;
			and/or influenza suspected	Callan et al., 1995; Early, 1992
1992	Bottlenose dolphins Tursiops truncatus (220)	(Calhoun and Aransas counties) Texas, U.S. Carbamates suspected	5. Carbamates suspected	Colbert et al., 1999; Duignan et al., 1996; Sweeney, 1992
1997–19	1992–1993 California sea lions Zalophus californianus (~1000) California 11S	)) California 11 S	El Niño 50 ginshot	Greio et al. 2005
1002	Houbon nouncion Phonons shows (21)	Moise Managhingth Compatibility	Eighouise interaction	Unlaw and Book 1002.
1995	Harbor porpoises <i>Procoena procoena</i> (64)	Maine, Massachusetts, Connecticut, New Jersey, Delaware, Virginia, North Carolina, U.S.	Fisheries interaction	Haley and Kead, 1993; Read and Murray, 2000
1993	Pinnipeds (53)(9 Steller Eumatopias jubatus;	Washington, U.S.	Gun shot. Bullets found on gross Norberg	Norberg
	15 California sea lions Zalophus californianus; 28 harbor seals Phoca vitulina; 1 unknown species sea lion)		necropsy and X-ray	[personal communication]
1994	Common dolphins Delphinus delphis	Black Sea	Morbillivirus	Birkun et al., 1999
1994	Common dolphins Delphinus delphis (53)	California, U.S.	Listed in Wilkinson, 1996, as a mortality event but not formally voted a UME by the Working Group. Etiology unknown	Reidarson et al., 1998

Table 1.	Continued			
Year	Species and no. affected	Location	Cause and comments	References
1994	Bottlenose dolphins Tursiops truncatus (72)	Texas, U.S.	Morbillivirus	Lipscomb et al., 1996
1995	California sea lions Zalophus californianus (222)	California, U.S.	Leptospirosis	Greig et al., 2005
1996	Sea otters Enhydra lutris (68)	(Cordova) Alaska, U.S.	Malnutrition, parasites, cold	Ballachey et al., 2002
1996	Right whales Eubalaena glacialis (6)	Florida, Georgia, U.S.	Blast injury suspected	Ridgway, 1996
1996	Florida manatees Trichechus manatus latirostris (149)	Florida (West coast), U.S.	Brevetoxin	Bossart et al., 1998; Landsberg and
				Steidinger, 1998
1996	Bottlenose dolphins Tursiops truncatus (30)	Mississippi, U.S.	Unknown. Coincident with algal bloom Working Group annual meeting	Working Group annual meeting
				report 1997 notes
1997	Mediterranean monk seals	Western Sahara	Harmful algal bloom/morbillivirus	Harwood, 1998; Hernandez et al.,
	Monachus monachus (150)			1998; Osterhaus et al., 1998
1997	Harbor seals Phoca vitulina (90)	California, U.S.	Pseudomonas aeruginosa associated	Working Group files, NMFS
			with pneumonia. Virus suspected.	
1998	Hooker's sea lions Phocarctos hookeri (60% of pups)	New Zealand	Unknown, bacteria likely	Baker, 1999
1998	California sea lions Zalophus californianus (70)	California, U.S.	Domoic acid	Scholin et al., 2000; Gulland,
				2000; Silvagni et al., 2005
1997–199	1997–1998 California sea lions Zalophus californianus (hundreds)	California, U.S.	El Niño	Greig et al., 2005
1999	Harbor porpoises Phocoena phocoena (216)	Maine, Massachusetts,	Oceanographic factors suggested	Marine Mammal Commission,
		Maryland, Virginia,	[Hohn, personal communication]	Annual Report to Congress
		North Carolina, U.S.		
2000	Caspian seals Phoca caspica (10,000)	Caspian Sea	Canine distemper virus (CDV)	Kennedy et al., 2000
2000	Beaked whales	Bahamas	Anthropogenic sound	Evans and England, 2001
2000	Sea otters Enhydra lutris (100)	(Cordova) Alaska, U.S.	Parasites ingested at a fish processing	Minutes of UME
			plant with discarded waste	Working Group meeting 2001
1999–200	1999–2000 Bottlenose dolphins Tursiops truncatus (115)	Florida (Panhandle), U.S.	Brevetoxin	Mase et al., 2000
1999–200	1999–2001 Gray whales Eschrichtius robustus (651)	California, Oregon, Washington, Alaska, U.S.; Canada; Mexico	Unknown. Starvation involved	Gulland et al., 2005; Moore et al., 2001
2000	California sea lions Zalophus californianus (178)	California, U.S.	Leptospirosis	Greig et al., 2005
2000	California sea lions Zalophus californianus (184)	California, U.S.	Domoic acid	Gulland et al., 2002
2000	Harbor seals <i>Phoca vitulina</i> (26)	California, U.S.	Unknown. Viral pneumonia suspected	Minutes of the UME Working Group meeting 2001, NMFS
				•

Cause and comments  Unknown.  Saxitoxin present in puffer fish Unknown  Malnutrition  Phocine distemper virus  Fle Domoic acid  Rebsiella pneumoniae  Cas embolism, sonar  Fle Brevetoxin  Domoic acid  Cas embolism, sonar  Fle Brevetoxin  Draceased detection,  voc ecological factors  Unknown. Saxitoxin and  3 humpbacks, respectively  To Chrenown  Brevetoxin  Unknown  Ha Leptospirosis	Table 1.	Continued			
Bottlenose dolphins Tursiops truncatus (35)  Hamp seals Phoca groenlandica (453)  Hawaiian monk seals Monachus schauinslandi (11)  Hawaiian lands, U.S.  Malnutrition  Hawaiian monk seals Monachus schauinslandi (11)  Hawaiian slands, U.S.  Multispecies(Common dolphins Delphinus delphis)  California sea lions Zalophus california delphis  Rocker's sea lion Phocarctos hookeri  Canary Islands Spain  Hooker's sea lion Phocarctos hookeri  California sea lions Zalophus californians delphis  California, U.S.  Domoic acid  Rocker's sea lion Phocarctos hookeri  Canary Islands Spain  California sea lions Zalophus californians delphis  Sea otters Enhydra lutris ~500)  Sea otters Enhydra lutris ~500  California, U.S.  Domoic acid detection,  California california californianus  sea otters Enhydra lutris ~500  Beluga whales Orlehimophus Californianus  Sea otters Enhydra lutris ~500  Beluga whales Orlehimophus Californianus  Sea otters Enhydra lutris ~500  Beluga whales Orlehimophus Californianus  Sea otters Enhydra lutris ~500  Beluga whales Orlehimophus Californianus  Sea otters Enhydra lutris ~500  Beluga whales Orlehimophus Californianus  Sea otters Enhydra lutris ~500  Beluga whales Orlehimophus Californianus  Sea otters Enhydra lutris ~500  Beluga whales Orlehimophus Californianus  Bottlenose dolphins Tursiops truncatus (107)  Bottlenose dolphins Tursiops truncatus (107)  Bottlenose dolphins California sea lions Zalophus californiaus (405)  California sea lions Zalophus californianus (405)  California sea lions Zaloph	Year	Species and no. affected	Location	Cause and comments	References
Harp seals Phoca groenlandica (453) Hawaiian Islands), U.S. Harbor seals Phoca vitulina (~25,000) Multispecies(Common dolphins Delphinus delphis; California, U.S. Harbor seals Phoca vitulina (~25,000) Multispecies(Common dolphins Delphinus delphis; California, U.S. Horida manates (14) Multispecies(Common dolphins Delphinus delphis; California, U.S. Horida manates Tridechus manatus lativastris (34) Florida (West coast), U.S. Beluga whales (14) Multispecies(Common dolphins Delphinus delphis; California, U.S. Beluga whales (15 humpback, 1 fin, 1 minks, 1 pilot, 2 unknown) Minks whales Balaenoptera acutorostrata Minks whales Balaenoptera acutorostrata Minks whales Balaenoptera acutorostrata Minks whales Balaenoptera acutorostrata  Exclusive Economic  Zone (EEZ)  Bottlenose dolphins Turisiops truncatus (107) Small cetaceans Small cetaceans California sea lions Zalophus californianus (405) Small cetaceans California sea lions Zalophus californianus (405) California, U.S. California, O.S. California, O.S. California, U.S. California, O.S. California, U.S. California,	2001	Bottlenose dolphins Tursiops truncatus (35)	Florida (Indian River Lagoon), U.S.	Unknown. Saxitoxin present in puffer fish	Barros, 2001 [unpublished]; Leighfield, 2002 [unpublished]
Hawaii (Northwest Mainetine scalations and (~25,000)  Hawaii (Northwest Hawaii (Northwest Hawaii (Northwest Hawaii (Northwest Harbor seals Phoca vitulina (~25,000)  Multispecies(Common dolphins Delphins delphis; California, U.S. Domoic acid detection, Sea otters Enhydra lurris (69)  Sea otters Enhydra lurris (69)  Sea otters Enhydra lurris (60)  Sea otters Enh	2001	Harp seals Phoca groenlandica (453)	Maine, Massachusetts, U.S.	•	Harris et al., 2002
Harbor seals Phoca vitulina (~25,000)  Multispecies(Common dolphins Delphinus delphis; California, U.S.  California sea lions Zalophus California, U.S.  Sea cotters Enhydra lutris; ~300)  Hooker's sea lion Phocarctos hookeri  Beaked whales (14)  Florida manatees Trichechus manatus lativostris (34) Florida (West coast), U.S.  Beluga whales (15)  Beluga whales Delphining Delphinus delphis; California, U.S.  California sea lions Zalophus California, U.S.  Beluga whales Delphining Latrices (20)  California vea lions Zalophus California, U.S.  Beluga whales Delphining Latrices (30)  California, U.S.  Large whales (16 humpback, 1 fin, 1 Maine, U.S.  Minke whales Balaenoptera acutorostrata  Exclusive Economic  Zone (EEZ)  Bottlenose dolphins Tursiops truncatus (107)  Small cetaceans (57)  Small cetaceans  Small cetaceans (57)  Small cetaceans  California, U.S.  Chikonia, U.S.  Brevetoxin and  Annice delega procere acutorostrata  Exclusive Economic  Zone (EEZ)  Bottlenose dolphins Tursiops truncatus (107)  Small cetaceans  Small cetaceans  Small cetaceans  California, U.S.  Unknown  Bas  Small cetaceans  Small cetaceans  Small cetaceans  Small cetaceans  California, U.S.  Unknown  Bas  Small cetaceans  California, U.S.  Unknown  Bas  Small cetaceans  S	2001	Hawanan monk seals Monachus schaumslandı (11)	Hawau (Northwest Hawaiian Islands), U.S.		Antonelis et al., 2001
Multispecies(Common dolphins Delphinus delphis; California, U.S. Domoic acid Maltispecies(Common dolphins Delphinus delphis; california sea lions Zalophus californianus; sea otters Ethydra lutris; ~500)  Hooker's sea lion Phocarctos hookeri Canary Islands, Spain Gas embolism, sonar Florende whales (14)  Horida manatees Trichechus manatus Intirostris (34) Florida (West coast), U.S. Domoic acid Maltispecies(Common dolphins Delphinus delphis; California, U.S. Domoic acid Maltispecies(Common dolphins Delphinus delphis; California, U.S. Domoic acid Gaetection, Sea otters Ethydra lutris (69)  Maine, U.S. Increased detection, Voluntown Saxitoria or Cook Inlet, Alaska, U.S. Increased detected in 2 and annihe, 1 pilot, 2 unknown)  Large whales (16 humpback, 1 fin, 1  Minke whales Balaenoptera acutorostrata  Exclusive Economic  Zone (EEZ)  Horida manatees Trichechus manatus lutriostris (96)  Small cetaceans (67)  Small cetaceans (67)  Small cetaceans (67)  Small cetaceans  California, Oregon, Leptospirosis Ra	2002	Harbor seals Phoca vitulina (~25,000)	Northern Europe; U.K.		Jensen et al., 2002
sea otters Enhydra lutris, ~500) Hooker's sea lion Phocarctos hookeri Beaked whales (14) Hooker's sea lion Phocarctos hookeri Canary Islands, Spain Garary Islands, Spain Gas embolism, sonar Florida divest coast), U.S. Brevetoxin Multispecies(Common dolphims Delphimus delphis; California, U.S. California, U.S. Sea otters Enhydra lutris (69) Sea otters Enhydra lutris (69) Beluga whales Delphimapteus leucas (20) Calok Inlet, Alaska, U.S. Large whales (16 humpback, 1 fin, 1 minke, 1 pilot, 2 unknown)  Large whales (16 humpback, 1 fin, 1 minke, 1 pilot, 2 unknown)  Minke whales Balaenoptena acutorostrata  Anninke whales Balaenoptena acutorostrata  Exclusive Economic  Zone (EEZ)  Florida (West coast), U.S. Brevetoxin  Bottlenose dolphins Tursiops truncatus (107)  Small cetaceans (67)  Small cetaceans  California, Oregon, California, Oregon, California, Oregon, California, Oregon, California, Oregon, California, Oregon, California, Spain  California, U.S. California, Oregon, California, Oregon, California, Oregon, California, Spain California, Oregon, California, Spain California, Oregon, California, Spain California, Spain California, Spain California, Spain California, Spain California, Spain California, California, Oregon, California, California, Spain California, California, Spain California, California, Spain California, California, California, Oregon, California, Califor	2002	Multispecies(Common dolphins Delphinus delphis; California sea lions Zalophus californianus;	California, U.S.		Mazet et al., 2005
Hooker's sea lion Phocarctos hookeri  Hooker's sea lion Phocarctos hookeri  Beaked whales (14)  Florida manatees Trichechus manatus latirostris (34)  Florida (West coast), U.S.  Domoic acid  California, U.S.  Domoic acid  Maine, U.S.  California, U.S.  California, U.S.  California, U.S.  California, U.S.  Domoic acid  Maine, U.S.  Cook Inlet, Alaska, U.S.  Unknown  Small cetaceans (57)  Small cetaceans  Codifornia, U.S.  Contriginia, U.S.  Colifornia, O.S.  Col		sea otters Enhydra lutris; $\sim$ 500)			
Beaked whales (14)  Florida manatees Trichechus manatus latirostris (34) Florida (West coast), U.S.  Multispecies(Common dolphins Delphinus delphis; California, U.S.  California sea lions Zalophus californiamus; sea otters Enlydra lutris; ~500)  Sea otters Enlydra lutris (69)  Beluga whales Delphinapteus leucas (20)  California, U.S.  California, U.S.  California, U.S.  Ecological factors  Domoic acid  Minke, U.S.  California, U.S.  Cook Inlet, Alaska, U.S	2002	Hooker's sea lion Phocarctos hookeri	New Zealand		Duignan et al., 2003
Florida manatees Trichechus manatus latirostris (34) Florida (West coast), U.S.  Multispecies(Common dolphins Delphinus delphis; California, U.S.  California sea lions Zalophus californianus; ~500)  Sea otters Enhydra lutris (99)  California, U.S.  Ecological factors  Domoic acid  Maine, U.S.  Ecological factors  Domoic acid  Maine, U.S.  Cook Inlet, Alaska, U.S.  Ecological factors  Unknown. Saxitoxin and  Janga whales (16 humpback, 1 fin, 1  Maine, U.S.  Maine, U.S.  Minke whales Balaenoptera acutorostrata  Exclusive Economic  Zone (EEZ)  Florida manatees Trichechus manatus latirostris (96) Florida (West coast), U.S.  Bottlenose dolphins Tursiops truncatus (107)  Small cetaceans (67)  Small cetaceans  California, U.S.  Unknown  Bassa otters Enhydra lutris  Brevetoxin  Minke whales Balaenoptera acutorostris (90)  Florida (Panhandle), U.S.  Unknown  Bassa otters Enhydra lutris  Brevetoxin  Minke whales Balaenoptera acutorostris (90)  Florida (Panhandle), U.S.  Unknown  Bassa California sea lions Zalophus californianus (405)  California, Oregon,  Leptospirosis  Represensin  Brevetoxin  Brevetoxin  Humbanatus Rairostris (405)  California, Oregon,  Leptospirosis	2002	Beaked whales (14)	Canary Islands, Spain		Jepson et al., 2003; Fernández et al., 2004
Multispecies(Common dolphins Delphinus delphis; California, U.S. Domoic acid M California sea lions Zalophus californianus; sea otters Enhydra lutris (69)  Sea otters Enhydra lutris (69) Sea otters Enhydra lutris (69) Sea otters Enhydra lutris (69) Sea otters Enhydra lutris (69) Cook Inlet, Alaska, U.S. Increased detection, V. ecological factors Beluga whales Delphinapteus leucas (20) Cook Inlet, Alaska, U.S. Increased detection, V. domoic acid detected in 2 and 3 humpback, respectively To minke, 1 pilot, 2 unknown)  Ainle, 1 pilot, 2 unknown)  Minke whales Balaenoptera acutorostrata  Exclusive Economic Zone (EEZ)  Florida manatees Trichechus manatus latirostris (96) Florida (West coast), U.S. Brevetoxin  Battlenose dolphins Tursiops truncatus (107)  Small cetaceans (67) Small cetaceans  California, U.S. Unknown  Humpbacks, respectively  California, U.S. Unknown  California sea lions Zalophus californiamus (405) California, Oregon, Leptospirosis	2002	Florida manatees Trichechus manatus latirostris (34)		Brevetoxin	Flewelling et al., 2005
California sea lions Zalophus californianus; sea otters Enhydra lutris (69) Sulforida manatees Phoca vitulina; Simal cetaceans (67) Small cetaceans (67) Small cetaceans (67) Small cetaceans Small cetaceans Small cetaceans California sea lions Zalophus californianus (405) California, Oregon, Lectospirosis Sea Ecological factors California Sea lions Zalophus californianus Sea otters Enhydra laticis Sea California sea lions Zalophus californianus Sea otters Enhydra laticis Sea otters Sea California Sea lions Salophus californianus Sea otters Sea California Sea lions Sea office Sea office Sea otters Sea otte	2003	Multispecies(Common dolphins Delphinus delphis;	California, U.S.		Minutes of the UMWE Working
sea otters Enhydra lutris, ~500) Sea otters Enhydra lutris (69) Cook Inlet, Alaska, U.S. Large whales Clé humpback, 1 fin, 1 Maine, U.S.  Cook Inlet, Alaska, U.S.  Unknown. Saxitoxin and a domoic acid detected in 2 and 3 humpbacks, respectively Some (BEZ)  Florida manatees Trichechus manatus latirostris (96) Florida (West coast), U.S.  Bottlenose dolphins Tursiops truncatus (107) Small cetaceans (67) Small cetaceans (67) Small cetaceans California sea lions Zalophus californianus (405) California, Oregon, Leptospirosis Ra		California sea lions Zalophus californianus;			Group annual meeting 2004
Sea otters Enhydra lutris (69) California, U.S. Beluga whales Delphinapteus leucas (20) Cook Inlet, Alaska, U.S. Increased detection, Vocok Inlet, Alaska, U.S. Large whales (16 humpback, 1 fin, 1) Maine, U.S.  Large whales (16 humpback, 1 fin, 1) Maine, U.S.  Large whales (16 humpback, 1 fin, 1) Maine, U.S.  Cook Inlet, Alaska, U.S.  Unknown. Saxitoxin and 3 humpbacks, respectively  Boulf of Maine, U.S.;  Unknown  To Alborov seals Phoca vitulina;  Minke whales Balaenoptera acutorostrata  Exclusive Economic  Zone (EEZ)  Horida manatees Trichechus manatus latirostris (96) Florida (West coast), U.S.  Bottlenose dolphins Tursiops truncatus (107)  Bottlenose dolphins Tursiops truncatus (107)  Small cetaceans (67)  Small cetaceans (67)  Small cetaceans  California sea lions Zalophus californianus (405)  California, Oregon,  Leptospirosis  Ecological factors  Vorginia, U.S.  Increased detection, Vorginia, U.S.  Increased detection, Vorginia, U.S.  Increased detection, Vorginia, U.S.  Unknown  Haborov Californianus (405)  California, Oregon,  Leptospirosis		sea otters Enhydra lutris; $\sim$ 500)			
Beluga whales Delphinapteus leucas (20)  Cook Inlet, Alaska, U.S.  Large whales (16 humpback, 1 fin, 1  minke, 1 pilot, 2 unknown)  Large whales (16 humpback, 1 fin, 1  minke, 1 pilot, 2 unknown)  Anumbacks, respectively  Coulf of Maine, U.S.;  Minke whales Balaenoptera acutorostrata  Exclusive Economic  Zone (EEZ)  Horida manatees Trichechus manatus latirostris (96) Florida (West coast), U.S.  Bottlenose dolphins Tursiops truncatus (107)  Bottlenose dolphins Tursiops truncatus (107)  Small cetaceans (67)  Small cetaceans (67)  Small cetaceans  California, U.S.  Unknown  Hinke whales Balaenopteus californianus (405)  California, Oregon,  Leptospirosis  Ra	2003	Sea otters Enhydra lutris (69)	California, U.S.		Draft report to UME Working Group, NMFS
Large whales (16 humpback, 1 fin, 1 Maine, U.S. Unknown. Saxitoxin and minke, 1 pilot, 2 unknown)  2004 Harbor seals Phoca vitulina;  Minke whales Balaenoptera acutorostrata Exclusive Economic  Zone (EEZ)  Florida manatees Trichechus manatus latirostris (96) Florida (West coast), U.S. Brevetoxin  Bottlenose dolphins Tursiops truncatus (107) Florida (Panhandle), U.S. Unknown  Small cetaceans (67) Virginia, U.S. Unknown  Small cetaceans  California sea lions Zalophus californianus (405) California, Oregon, Leptospirosis	2003	Beluga whales Delphinapteus leucas (20)	Cook Inlet, Alaska, U.S.		Vos and Shelden, 2005
Large whales (16 humpback, 1 fin, 1  minke, 1 pilot, 2 unknown)  -2004 Harbor seals Phoca vitulina;  Minke whales Balaenoptera acutorostrata  Minke whales Balaenoptera acutorostrata  Exclusive Economic  Zone (EEZ)  Florida manatees Trichechus manatus latirostris (96)  Bottlenose dolphins Tursiops truncatus (107)  Small cetaceans (67)  Small cetaceans  California sea lions Zalophus californianus (405)  California, Oregon,  Leptospirosis  Unknown  Unknown  Harbor saxitoxin and Day  Bomoic acid detected in 2 and  3 humpbacks, respectively  Transmit and 2 and  Bordle of Maine, U.S.;  Unknown  Chikown  California, Oregon,  Leptospirosis  Exclusive Economic  Small cetaceans  California, Oregon,  Leptospirosis				ecological factors	
minke, 1 pilot, 2 unknown)  2004 Harbor seals Phoca vitulina;  Minke whales Balaenoptera acutorostrata  Exclusive Economic  Zone (EEZ)  Florida manatees Trichechus manatus latirostris (96) Florida (West coast), U.S.  Bottlenose dolphins Tursiops truncatus (107)  Small cetaceans (67)  Small cetaceans  California sea lions Zalophus californianus (405)  California, Oregon,  Leptospirosis  domoic acid detected in 2 and 3 humpbacks, respectively Transpace of the call of Maine, U.S.  Brevetoxin  Minke whales Balaenoptera acutorostrata  Rabinandes, U.S.  Unknown  High Carolina, U.S.  California, Oregon,  Leptospirosis  Rabinandes  California vea lions Zalophus californianus (405)  California, Oregon,  Leptospirosis	2003	Large whales (16 humpback, 1 fin, 1	Maine, U.S.		Draft report to UME Working Group, NMFS
3 humpbacks, respectively  2004 Harbor seals Phoca vitulina; Minke whales Balaenoptera acutorostrata  Exclusive Economic Zone (EEZ)  Florida manatees Trichechus manatus latirostris (96) Florida (West coast), U.S.  Bottlenose dolphins Tursiops truncatus (107)  Small cetaceans (67)  Small cetaceans  California sea lions Zalophus californianus (405)  California, U.S.  Unknown  Harbor sepectively  Torknown  Brevetoxin  Dy North Carolina, U.S.  Unknown  Harbor sealions Zalophus californianus (405)  California, Oregon,  Leptospirosis		minke, 1 pilot, 2 unknown)		domoic acid detected in 2 and	
Alinke whales Balaenoptera acutorostrata Exclusive Economic  Minke whales Balaenoptera acutorostrata  Zone (EEZ)  Florida manatees Trichechus manatus latirostris (96) Florida (West coast), U.S. Brevetoxin  Bottlenose dolphins Tursiops truncatus (107)  Small cetaceans (67)  Small cetaceans (67)  Small cetaceans  California sea lions Zalophus californianus (405)  California, Oregon,  Leptospirosis  Trichochus  Brevetoxin  Dry  Worth Carolina, U.S.  Unknown  Ha  Ra				3 humpbacks, respectively	
Minke whales Balaenoptera acutorostrata  Zone (EEZ)  Florida manatees Trichechus manatus latirostris (96) Florida (West coast), U.S. Brevetoxin  Bottlenose dolphins Tursiops truncatus (107)  Small cetaceans (67)  Small cetaceans  California sea lions Zalophus californianus (405)  California, Oregon,  Leptospirosis  Ra	2003-2004	Harbor seals Phoca vitulina;	Gulf of Maine, U.S.;	Unknown	Touhey [personal communication]
Zone (EEZ) Florida manatees Trichechus manatus latirostris (96) Florida (West coast), U.S. Brevetoxin  Bottlenose dolphins Tursiops truncatus (107) Florida (Panhandle), U.S. Brevetoxin  Small cetaceans (67) Virginia, U.S. Unknown  Small cetaceans  California sea lions Zalophus californianus (405) California, Oregon, Leptospirosis Re		Minke whales Balaenoptera acutorostrata	Exclusive Economic		
Florida manatees Trichechus manatus latirostris (96) Florida (West coast), U.S. Brevetoxin  Bottlenose dolphins Tursiops truncatus (107) Florida (Panhandle), U.S. Brevetoxin  Small cetaceans (67) Virginia, U.S. Unknown  Small cetaceans  California sea lions Zalophus californianus (405) California, Oregon, Leptospirosis Ra			Zone (EEZ)		
Bottlenose dolphins Tursiops truncatus (107) Florida (Panhandle), U.S. Brevetoxin Dr Small cetaceans (67) Virginia, U.S. Unknown Hamble California sea lions Zalophus californianus (405) California, Oregon, Leptospirosis Ra	2003	Florida manatees Trichechus manatus latirostris (96)		Brevetoxin	Minutes of the UME Working Group annual meeting 2004
Small cetaceans (67)  Small cetaceans  Small cetaceans  California sea lions Zalophus californianus (405)  Virginia, U.S.  Unknown  Horizonia, Oregon,  Leptospirosis	2004	Bottlenose dolphins Tursiops truncatus (107)	Florida (Panhandle), U.S.		Draft report to UME Working Group;
Small cetaceans (67)  Small cetaceans  Small cetaceans  North Carolina, U.S.  Unknown  California sea lions Zalophus californianus (405)  California, Oregon,  Leptospirosis					Gaydos [in preparation]; Flewelling et al., 2005
Small cetaceans North Carolina, U.S. Unknown California sea lions <i>Zalophus californianus</i> (405) California, Oregon, Leptospirosis	2004	Small cetaceans (67)	Virginia, U.S.	Unknown	Barco, report to Working Group
California sea lions Zalophus californianus (405) California, Oregon, Leptospirosis	2004	Small cetaceans	North Carolina, U.S.		Hohn [personal communication]
	2004	California sea lions Zalophus californianus (405)	California, Oregon,		Raverty et al., 2005
Washington, U.S.; Canada			Washington, U.S.; Canada		

Table 1. Continued	ntinued			
Year	Species and no. affected	Location	Cause and comments	References
2005	Manatees Trichechus manatus; Bottlenose dolphins Tursiops truncatus (ongoing Dec. 2005)	Florida (West coast), U.S.	Brevetoxin. Bird, turtle, and fish kills associated with the event	
2005	Harbor porpoises  Phocoena phocoena (ongoing Dec. 2005)	North Carolina, U.S.	Unknown	Hohn [personal communication]
2005	California sea lions Zalophus californianus; Northern fur seals Callorhinus ursinus (several hundred)	California, U.S.	Domoic acid	Goldstein et al., 2005
2005 2005 2005–2006	Mixed cetaceans Large whales Bottlenose dolphins <i>Tursiops truncatus</i>	North Carolina, U.S. Eastern North Atlantic, U.S. Florida, U.S.	Unknown Domoic acid suspected Brevetoxin suspected	Hohn et al., 2006 Touhey [personal communication]

<sup>a</sup>Data are obtained from the published literature and the National Marine Fisheries Service database

## DISCUSSION AND CONCLUSIONS

This survey of the peer reviewed literature on marine mammal diseases and reports of marine mammal mass mortality events suggests that there has indeed been an increase in the frequency of marine mammal die-offs resulting from exposure to harmful algal blooms over the past 40 years. The evidence for changes in frequency of other types of diseases is difficult to interpret, particularly in recent years for a number of reasons. These include changes in investigation effort, improved methodology for detection of disease, biases associated with sources of animals for detection of disease, and changes in the population density of some coastal marine mammal populations that are more easily studied.

Effort aimed at identifying disease has changed due to the underlying increasing global trend in the reporting of marine mammal disease in the literature and in the documenting of mass mortalities associated with a general increase in scientific research and in the number of researchers in this field. This latter increase was previously noted by Lavigne et al. (1999) and attributed to changing attitudes to marine mammals. For example, the membership of the Society for Marine Mammalogy, since its inception in 1981 to its most recent conference in 2005, has increased approximately threefold from  $\sim$ 780 to  $\sim$ 2300.

Over the last 40 years, many marine mammal populations have increased significantly, due to changes in hunting and harvesting pressure and possibly to changes in food availability where ecosystem regime changes have occurred. This not only will influence the availability of carcasses for detection, but can allow density-dependent diseases to increase in prevalence. If a certain proportion of a population dies due to environmental changes and further selective pressures result in malnutrition, large increases in marine mammal populations may result in clusters of malnourished animals dying, increasing the likelihood of their detection by humans. Mass mortality events associated with starvation are reviewed in Geraci et al. (1999). Density-dependent diseases will increase as marine mammal populations recover. Epidemiological models of PDV suggest the population of harbor seals in Europe is not large enough to sustain the disease as endemic, epidemics only occurring when the pool of susceptible seals is over a critical threshold (Grenfell et al., 1992) and the disease will fade out between outbreaks. The first epidemic of PDV in European harbor seals in 1988-1989 was unexpected and required considerable effort to identify its etiology (Kennedy et al., 1988). However, disease modeling exercises predicted that another epidemic was likely after approximately 10 years based on knowledge of morbillivirus epidemiology and harbor seal population dynamics (Grenfell et al., 1992). Another epidemic of PDV occurred in European harbor seals in 2002, and although many details of the epidemic are still unclear (Harkonen et al., 2006), these recurring epidemics at approximately a decade interval do not suggest a change in harbor seal health over this period, but reflect changes in population immunity. A much longer dataset (i.e., 100 years) would be needed to detect changes in frequency of morbillivirus epidemics in seals at this interepidemic interval. Nothing is known about the diseases that may have been introduced into many marine mammal populations prior to 1960, but which did not establish themselves because populations were historically too small for the disease to propagate between individuals.

The change in methodology used to study marine mammal health and disease can also influence the frequency of novel disease detection. The change from studying harvested and by-caught animals to studying stranded and rehabilitated animals will result in a higher detection of diseased animals. An increase in the numbers of individuals stranding in a location can indicate an increase in mortality rate (Heide-Jorgensen et al., 1992; Harkonen et al., 2006), although stranding rates and numbers often do not correlate directly with mortality rates due to a variety of factors influencing carcass deposition and detection (Eguchi, 2002; Thompson et al., 2005). Documentation of stranding events has improved globally over the last 40 years, with the earliest organized attempts originating in the United Kingdom (U.K.) (Fraser, 1934, 1946, 1953, 1956; Geraci, 1978; Sergeant, 1982; Dierauf and Gulland, 2001).

Stranded animals are a useful source of information on diseases occurring in marine mammals, as they are more readily available to pathologists than free-ranging animals. A number of emerging disease entities, suites of lesions and infectious agents in marine mammals were first identified in stranded animals, after which their presence in the freeranging population was confirmed. Pathogens identified through stranding programs include phocine distemper virus (PDV) that caused the death of over 18,000 harbor (Phoca vitulina) seals in Europe in 1988 and over 20,000 in 2002 (Osterhaus et al., 1988; Kennedy et al., 1988; Jensen et al., 2002; Harkonen et al., 2006), phocine herpesvirus (PhHV1) isolated from stranded harbor seals in 1985

(Osterhaus et al., 1985), and Brucella marinus in a variety of species (Ross et al., 1996; Garner et al., 1997). Stranded animals can also alert us to diseases that are present in the more inaccessible wild animals that would be difficult to detect in random samplings of such populations. For example, 17% of sexually mature California sea lions (Zalophus californianus) that stranded and died along the northern coast of California showed neoplasia when examined postmortem (Gulland et al., 1996b). In comparison, only one case of neoplasia has been observed in California sea lions at rookeries on San Miguel Island (CA) where there are more than 100,000 sea lions [Spraker, personal communication]. The study of neoplasia pathogenesis is thus more readily performed on stranded sea lions than on those in rookeries, and stranded animals essentially serve as sentinels for their wild con-specifics. Animals in rehabilitation also provide a highly significant sampling method for the study of marine mammal disease. Indeed, the discovery that domoic acid produced by harmful algal blooms of Pseudo-nitzschia australis was causing mortality and seizures in California sea lions was entirely as a result of a comprehensive rescue and rehabilitation program for the affected region (Scholin et al., 2000).

Stranded animals, however, do not constitute an ideal system for the study of disease, as they do not represent the entire population (Aguilar and Borrell, 1994). In addition, samples of stranded animals are rarely age and sex structured, and biological data such as individual life histories, feeding habits, reproductive success, or disease progression are not typically available. Examination of both live and dead strandings are useful for detecting disease but are limited when the objective is to assess the prevalence of a disease and its impact on a host population, since the sample will be skewed towards those likely to come ashore. Decomposition is also a problem for obtaining an accurate diagnosis and good histology among dead strandings, while human concern and intervention can interfere with research on live stranded animals.

The examination of stranded cetaceans has also allowed the detection of a novel suite of lesions, termed "gas bubble disease" (Jepson et al., 2003). Animals with these lesions stranded in association with navy exercises, but whether the frequency of mortality of cetaceans from these lesions is changing over time in association with increasing levels of ocean noise cannot be determined due to the lack of a systematic time series on pathology of stranded cetaceans worldwide. The few case studies on this syndrome are a result of the fortuitous proximity of skilled veterinary pathologists, fresh stranded cetaceans, and publicly accessible information on naval activities. To determine the extent of this problem and whether its frequency is changing, improved facilities and logistic support for examining dead marine mammals thoroughly are needed along coastlines throughout the world.

It is possible that prior to the recent interest in ocean health, some marine mammal disease outbreaks may have been missed. However, high numbers of dead animals are difficult to ignore—there have been reports of seals suffering "epizooty" in the literature from the 1700s (see Harwood and Hall, 1990)—so it is unlikely that epidemics would have been missed prior to the first of the recent events in the 1980s. Events are more likely to be documented in the peer-reviewed literature when the causes were identified and can be acknowledged retrospectively, than events with no identified causes. Thus, there may have been a number of marine mammal mortality events of unknown etiology between 1978 and 1992 that are not recorded in the published literature reviewed here. As technology advances, so have methods to detect toxins and pathogens from smaller samples of less well-preserved tissues. This has resulted in the changes in frequency that etiological agents have been described in the published literature (Fig. 2). There is also variability worldwide in the availability of more advanced laboratory techniques for examining marine mammal tissues, as well as a lack of diagnostic tests validated for marine mammals. To prevent this bias from continuing to impair our ability to detect changes in marine mammal health, dedicated laboratories for marine mammals are needed.

To determine whether marine mammal health is indeed changing, a concerted effort is needed worldwide to coordinate investigations into marine mammal die-offs, to investigate associations between disease and ecological variables, to share data on methodology used for investigations and laboratory studies, and to document the stranding response effort. Over time, we will be able to use geographic information systems (GIS) and spatial epidemiological studies to investigate temporal trends in marine mammal disease. In 1988, the first dramatic PDV epidemic that killed over 18,000 harbor seals in Europe raised awareness of the need for contingency plans to investigate marine mammal die-offs, and for long-term monitoring of strandings (Heide-Jorgensen et al., 1992; Thompson and Hall, 1993). In 1989, the Department of the Environment (now the Department for the Environment, Food and

Rural Affairs) in the U.K. established a national program to investigate marine mammal mortalities, and to coordinate responses. In the U.S., three specific events triggered the development of a legal framework that addressed marine mammal unusual mortality events. The first was a stranding of 14 humpback whales (Megaptera novaeangliae) off Cape Cod, Massachusetts, in 1987 (Geraci et al., 1989). The second was a bottlenose dolphin die-off along the Atlantic seaboard between 1987 and 1988 (Geraci, 1989), and the third the Exxon Valdez oil spill in Prince William Sound, Alaska, in 1989 (Loughlin, 1994). As a response to the increased concerns over these die-offs and the potential for environmental degradation, the Marine Mammal Health and Stranding Response Act was established, and became Title IV of the Marine Mammal Protection Act (MMPA) with a provision for responses to marine mammal unusual mortality events. There are four aims of these responses: (1) to minimize death and suffering, (2) to determine the cause of the event, (3) to determine the effect of the event on the population, and (4) to identify the role of environmental variables in the event; only the first two of these have been significantly addressed since the UME program was established in 1992. Efforts to minimize death and suffering have improved as an increasing number of stranding network facilities employ veterinarians to advise on medical care of stranded marine mammals, and the field of marine mammal medicine has developed considerably over the last 10 years. Identification of the cause of mortality events has been achieved in just over half (56%) of the investigations, some of these resulting in the detection of novel disease syndromes in marine mammals (Geraci et al., 1989; Lipscomb et al., 1994; Bossart et al., 1998; Scholin et al., 2000).

Identification of novel diseases requires significant resources and collaboration, thus achieving this through investigations of mortality of wild mammals is a considerable accomplishment. However, trends in the frequency of these diseases, the impact of environmental change on these diseases, and the effects of these mortality events on host populations cannot be determined from the data currently available. The available data are adequate to identify an increase in frequency of marine mammal die-offs in the United States due to poisoning by harmful algal blooms, but other changes in marine mammal health worldwide cannot be determined in this study. Without this additional information, the status of marine mammal health and the impacts of anthropogenic activities and environmental factors, such as climate change, cannot be assessed. To improve our understanding of marine mammal health and

determine the impacts of human activities on these animals, improved programs for disease surveillance, emergency response, laboratory diagnosis, and research into contributing factors are needed, with international collaborations to address marine mammal health in ocean ecosystems without national boundaries.

### **A**CKNOWLEDGMENTS

We particularly thank Dr. Teri Rowles, NOAA, National Marine Fisheries Service (NMFS), Center for Marine Animal Health, for her encouragement and support. We also thank all members of the National Marine Mammal Health and Stranding Network for their hard work in collecting the data presented here and caring for the stranded animals, and the staff of the NMFS for managing the Marine Mammal Health and Stranding Program, especially Janet Whaley, Trevor Spradlin, Sarah Wilkin, Michelle Ordono, Angela Collins-Payne, Joe Cordaro, and Brent Norberg. We also thank Denise Greig for assistance with this manuscript.

#### REFERENCES

- Aguilar A, Borrell A (1994) Abnormally high polychlorinated biphenyl levels in striped dolphins (*Stenella coeruleoalba*) affected by the 1990–1992 Mediterranean epizootic. *Science of the Total Environment* 154:237–247
- Aguilar A, Raga JA (1993) The striped dolphin epizootic in the Mediterranean Sea. *Ambio* 22:524–528
- Antonelis GB, Ryon R, Braun R, Spraker T, Baker J, Rowles T (2001) Juvenile Hawaiian monk seal (Monachus schauinslandi) unusual mortality event in the northwestern Hawaiian Islands. In: Society for Marine Mammalogy, Proceedings of the 14th Biennial Conference on the Biology of Marine Mammals, Vancouver, Canada, pp 7–8 (abstract)
- Baker A (1999) Unusual Mortality of the New Zealand Sea Lion, Phocarctos hookeri, Auckland Islands, January–February 1998.Department of Conservation, Wellington, New Zealand, 84 pp
- Ballachey BE, Gorbics CS, Doroff AM (2002) Sea otter mortality in Orca Inlet, Prince William Sound, Alaska, winter 1995–1996. USFWS Technical Report MMM 02-1, 26 pp
- Barros N (2001) Preliminary analysis of stomach contents of bottlenose dolphins from the Indian River Lagoon unusual mortality event, June–July 2001. Unpublished report to the National Marine Fisheries Service, 5 pp
- Birkun A, Kuiken T, Krivokhizhin S, Haines DM, Osterhaus ADME, van de Bildt MWG, et al. (1999) Epizootic of morbilliviral disease in common dolphins (*Delphinus delphis ponticus*) from the Black Sea. *Veterinary Record* 144:85–92
- Bossart GD, Baden DG, Ewing RY, Roberts B, Wright S (1998) Brevetoxicosis in manatees (*Trichechus manatus latirostris*) from

- the 1996 epizootic: gross, histologic and immunohistologic features. *Toxicologic Pathology* 26:276–282
- Callan RJ, Early G, Kida H, Hinshaw VS (1995) The appearance of H3N3 influenza viruses in seals. *Journal of General Virology* 76:199–203
- Colbert AA, Scott GI, Fulton MH, Wirth EF, Daugomah JW, Key PB, et al. (1999) Investigation of unusual mortalities of bottlenose dolphins along the mid-Texas coastal bay ecosystem during 1992. U.S. Department of Commerce, Seattle, WA: NOAA Technical Report NMFS 147, 23 pp
- Cox TM, Ragen TJ, Read AJ, Vos E, Baird RW, Balcomb K, et al. (2006) Understanding the impacts of anthropogenic sound on beaked whales. *Journal of Cetacean Research and Management* 7:177–187
- Cunningham A (2005) A walk on the wild side—emerging wildlife diseases. *British Medical Journal* 331:1214–1215
- DeGange AR, Vacca M (1989) Sea otter mortality at Kodiak Island, Alaska, during summer 1987. *Journal of Mammalogy* 70:836–838
- DeSwart RL, Ross PS, Vedder LJ, Timmerman HH, Heisterkamp S, Van Loveren H, et al. (1994) Impairment of immune function in harbor seals (*Phoca vitulina*) feeding on fish from polluted waters. *Ambio* 23:155–159
- Dierauf L, Gulland FMD (2001) Handbook of Marine Mammal Medicine, 2nd ed. Boca Raton, FL: CRC Press, 1063 pp
- Dierauf L, Vandenbroek DJ, Roletto J, Koski M, Amaya L, Gage LG (1985) An epizootic of leptospirosis in California sea lions. Journal of the American Veterinary Medical Association 187:1145–1148
- Domingo M, Visa J, Pumarola M, Marco AJ, Ferrer L, Rabanal R, et al. (1992) Pathological and immunocytochemical studies of morbillivirus infection in striped dolphins (*Stenella coeruleoalba*). Veterinary Pathology 29:1–10
- Duignan PJ, House C, Odell DK, Wells RS, Hansen LJ, Walsh MT, et al. (1996) Morbillivirus infection in bottlenose dolphins: evidence for recurrent epizootics in the western Atlantic and Gulf of Mexico. *Marine Mammal Science* 12:499–515
- Duignan PJ, Wilkinson I, Alley MR (2003) New Zealand sea lion (*Phocarctos hookeri*) epidemic 2002. New Zealand Veterinary Journal 51:46
- Early G (1992) Summary Report and Evaluation of the 1992 Seal Distemper Testing Program with Comparison of Mortality Rates. In fulfillment of requisition number: NFFM1020200394
- Eguchi T (2002) A method for calculating the effect of a die-off from stranding data. *Marine Mammal Science* 18:698–709
- Epstein P (1996) Emergent stressors and public health implications in large marine ecosystems: an overview. In: Smayda T (editor), Northeastern Shelf Ecosystem: Assessment, Sustainability, and Management Cambridge, MA: Blackwell, pp 417–38
- Evans DL, England GR (editors) (2001) Joint Interim Report Bahamas Marine Mammal Stranding Event of 14–16 March 2000. National Oceanic and Atmospheric Administration/US Navy, 61 pp. Available: http://www.nmfs.noaa.gov/prot\_res/ overview/Interim\_Bahamas\_Report.pdf
- Fernández A, Arbelo M, Deaville R, Patterson IAP, Castro P, Baker JR, et al. (2004) Beaked whales, sonar and decompression sickness. *Nature* 10:1038
- Flewelling LJ, Naar JP, Abbott JP, Baden DG, Barros NB, Bossart GD, et al. (2005) Red tides and marine mammal mortalities. *Nature* 435:755–756

- Fraser FC (1934) Report on cetacea stranded on the British coast from 1927-1932. British Museum of Natural History 11
- Fraser FC (1946) Report on cetacea stranded on the British coast from1933–1937. British Museum of Natural History 12
- Fraser FC (1953) Report on cetacea stranded on the British coast from 1938-1947. British Museum of Natural History 13
- Fraser FC (1956) Report on cetacea stranded on the British coast from 1948-1956. British Museum of Natural History 14
- Friedlaender AS (2000) Using strandings for effective management and conservation of bottlenose dolphins (Tursiops truncatus) along the U.S. Atlantic coast. MS Thesis. Department of Biological Sciences, University of North Carolina at Wilmington, 85 pp
- Gardner TA, Cote IM, Gill JA, Grant A, Watkinson AR (2003) Long-term region-wide declines in Caribbean corals. Science 301:958-960
- Garner MM, Lambourn DM, Jeffries SJ, Hall PB, Rhyan JC, Ewalt DR, et al. (1997) Evidence of Brucella infection in Parafilaroides lungworms in a Pacific harbor seal (Phoca vitulina richardsi).. Journal of Veterinary Diagnostic Investigation 9:298-303
- Geraci JR (1978) The enigma of marine mammal strandings. Oceanus 21:38-47
- Geraci JR (1989) Clinical investigation of the 1987-1988 mass mortality of bottlenose dolphins along the US central and south Atlantic coast. Washington, DC: Final Report, US Marine Mammal Commission, 63 pp
- Geraci JR, Anderson DM, Timperi RJ, St. Aubin DJ, Early GA, Prescott JH, et al. (1989) Humpback whales (Megaptera novaeangliae) fatally poisoned by dinoflagellate toxin. Canadian Journal of Fisheries and Aquatic Sciences 46:1895-1898
- Geraci JG, Duignan P, Early G (1993) Survey for morbillivirus in pinnipeds along the northeastern coast. Final Report to NOAA/ NMFS, Contract No. 50-DGNF-2-00098
- Geraci JR, Harwood J, Lounsbury VJ (1999) Marine mammal dieoffs. In: Twiss JR, Reeves RR (editors), Conservation and Management of Marine Mammals Washington, DC: Smithsonian Institution Press, pp 367-395
- Geraci JR, St. Aubin DJ, Barker IK, Webster RG, Hinshaw VS, Bean WJ, et al. (1982) Mass mortality of harbor seals: pneumonia associated with influenza A virus. Science 215:1129-1131
- Gilmartin WG, DeLong RL, Smith AW, Griner LA, Dailey MD (1980) An investigation into unusual mortality in the Hawaiian monk seal, Monachus schauinslandi. In: Proceedings of the Symposium on Status of Resource Investigation in the Northwestern Hawaiian Island, April 24-25, 1980, Grigg RW, Pfund RT (editors), University of Hawaii, Sea Grant Rep. UNIHI-SEAGRANT-MR-80-04, pp 32-41
- Goldstein T, Gulland F, Langlois G, Silver M, Haulena M, Lowenstine LJ, et al. (2005) Sub-lethal and long term effects of exposure to domoic acid in stranded California sea lions (Zalophus californianus). In: Proceedings of the 16th Biennial Conference on the Biology of Marine Mammals, San Diego, CA, December 12–16, 2005, p 109
- Grachev MA, Kumarev VP, Mamaev LV, Zorin VL, Baranova LV, Denikina NN, et al. (1989) Distemper virus in Baikal seals. Nature 338:209-210
- Greig DJ, Gulland FMD, Kreuder C (2005) A decade of live California sea lion (Zalophus californianus) strandings along the central California coast: causes and trends, 1991-2000. Aquatic Mammals 31:40-51

- Grenfell BT, Lonergan ME, Harwood J (1992) Quantitative investigations of the epidemiology of phocine distemper virus (PDV) in European common seal populations. Science of the Total Environment 115:15-29
- Gulland F (2000) Domoic acid toxicity in California sea lions (Zalophus californianus) stranded along the central California coast, May-October 1998. NOAA Technical Memorandum, NMFS-OPR, 17, 45 pp
- Gulland F, Pérez-Cortés H, Urbán MJ, Rojas-Bracho RL, Ylitalo G, Weir J, et al. (2005) Eastern North Pacific gray whale (Eschrichtius robustus) unusual mortality event, 1999-2000: a compilation. U.S. Department of Commerce, NOAA Technical Memorandum, NMFS-AFSC-150, 33 p
- Gulland FMD, Haulena M, Fauquier D, Langlois G, Lander ME, Zabka T, et al. (2002) Domoic acid toxicity in Californian sea lions (Zalophus californianus): clinical signs, treatment and survival. Veterinary Record 150:475-480
- Gulland FMD, Koski M, Lowenstine LJ, Colagrass A, Morgan L, Spraker T (1996a) Leptospirosis in California sea lions (Zalophus californianus) stranded along the central California coast, 1981-1994. Journal of Wildlife Diseases 32:572-580
- Gulland FMD, Trupkiewicz JG, Spraker TR, Lowenstine LJ (1996b) Metastatic carcinoma of probable transitional cell origin in free-living California sea lions (Zalophus californianus): 64 Cases (1979-1994). Journal of Wildlife Diseases 32:250-258
- Haley NJ, Read A (1993) Summary of the workshop on harbor porpoise mortalities and human interactions. NOAA Technical Memorandum NMFS-F/NER-5, 32 pp
- Hall AJ, Law RJ, Wells DE, Harwood J, Ross HM, Kennedy S, et al. (1992) Organochlorine levels in common seals (Phoca vitulina) that were victims and survivors of the 1988 phocine distemper epizootic. Science of the Total Environment 115:145-162
- Hammond JA, Hall AJ, Dyrynda L (2005) Comparison of polychlorinated biphenyl (PCB) induced effects on innate immune functions in harbour and grey seals. Aquatic Toxicology 74:126-
- Hansen LJ (1992) Report on investigation of 1990 Gulf of Mexico bottlenose dolphin strandings. NOAA-NMFS-SEFSC Contribution: MIA-92/93-21. 219 pp
- Harkonen T, Dietz R, Reijnders PJH, Teilmann J, Harding K, Hall AJ, et al. (2006) A review of the 1988 and 2002 phocine distemper virus epidemics in European harbour seals. Diseases of Aquatic Organisms 68:115-130
- Harris DE, Lelli B, Jakush G (2002) Harp seal records from the southern Gulf of Maine: 1997-2001. Northeastern Naturalist 9:331-340
- Harvell CD, Kim K, Burkholder J, Colwell RR, Epstein PR, Grimes J, et al. (1999) Emerging marine diseases—climate links and anthropogenic factors. Science 285:1505-1510
- Harvell CD, Mitchell CE, Ward JR, Altizer S, Dobson AP, Ostfield RS, et al. (2002) Climate warming and disease risks for terrestrial and marine biota. Science 296:2158-2162
- Harvell D, Aronson RA, Baron N, Connell J, Dobson A, Ellner S, et al. (2004) The rising tide of ocean diseases: unsolved problems and research priorities. Frontiers in Ecology and Environment 2:375-382
- Harwood J (1998) What killed the monk seals?. Nature 393:17–18
- Harwood J, Hall A (1990) Mass mortality in marine mammals: its implications for population dynamics and genetics. Trends in Ecology and Evolution 5:254-257

- Heide-Jorgensen MP, Harkonen T, Dietz R, Thompson PM (1992) Retrospective of the 1988 European seal epizootic. Diseases of Aquatic Organisms 13:37–62
- Hernandez M, Robinson I, Aguilar A, Gonzalea LM, Lopez-Jurado LF, Reyero MI, et al. (1998) Did algal toxins cause monk seal mortality?. *Nature* 393:28–29
- Hinshaw VS, Bean WJ, Webster RG, Rehg JE, Fiorelli P, Early G, et al. (1984) Are seals frequently infected with avian influenza viruses?. *Journal of Virology* 51:863–865
- Hohn AH, Rotstein DR, Harms CA, Southall BL (2006) Report on marine mammal unusual mortality event UMESE0501Sp: multispecies mass stranding of pilot whales (*Globicephala macrorhynchus*), minke whale (*Balaenoptera acutorostrata*), and dwarf sperm whales (*Kogia sima*) in North Carolina on 15–16 January 2005. U.S. Department of Commerce National Oceanic and Atmospheric Administration, National Marine Fisheries Service, NOAA Technical Memorandum NMFS-SEFSC-537, March 2006
- Jensen T, van de Bildt M, Dietz H, Andersen T, Hammer A, Kuiken T, et al. (2002) Another phocine distemper outbreak in Europe. *Science* 297:209
- Jepson PD, Arbelo M, Deaville R, Patterson IAP, Castro P, Baker JR, et al. (2003) Gas-bubble lesions in stranded cetaceans. *Nature* 425:575–576
- Kennedy S, Kuiken T, Jepson P, Deaville R, Forsyth M, Barrett T, et al. (2000) Mass die-off of Caspian seals caused by canine distemper virus. *Emerging Infectious Diseases* 6:637–639
- Kennedy S, Smyth JA, McCullough SJ, Allan GM, McNeilly F, McQuaid S (1988) Confirmation of cause of recent seal deaths. *Nature* 335:404
- Kim K, Dobson AP, Gulland FMD (2005) Diseases and the conservation of marine biodiversity. In: Norse EA, Crowder LB (editors), Marine Conservation Biology: The Science of Maintaining the Sea's Biodiversity Washington, DC: Island Press, pp 149–166
- Knap A, Dewailly E, Furgal C, Galvin CJ, Baden D, Bowen RE, et al. (2002) Indicators of ocean health and human health: developing a research and monitoring framework. *Environmental Health Perspectives* 110:839–845
- Knowlton N (2004) Ocean health and human health: developing a research monitoring framework. *Environmental Health Perspectives* 112:A262
- Kuehl DW, Haebler R (1995) Organochlorine, organobromine, metal, and selenium residues in bottlenose dolphins (*Tursiops truncatus*) collected during an unusual mortality event in the Gulf of Mexico, 1990. *Archives of Environmental Contaminants and Toxicology* 28:494–499
- Kuiken T, Simpson VR, Allchin CR, Bennett PM, Codd GA, Harris EA, et al. (1994) Mass mortality of common dolphins (*Delphinus delphis*) in south west England due to incidental capture in fishing gear. *Veterinary Record* 134:81–89
- Lafferty KD, Porter J, Ford SE (2004) Are diseases increasing in the ocean?. *Annual Review of Ecological Systems* 35:31–54
- Landsberg JH, Steidinger KA (1998) A historical review of *Gymnodinium breve* red tides implicated in mass mortalities of the manatee (*Trichechus manatus latirostris*) in Florida, USA. In: *Harmful Algae*, Reguera B, Blanco J, Fernández ML, Wyatt T (editors), Paris: Xunta de Galicia, IOC, pp 97–100
- Lavigne DM, Scheffer VB, Kellert SR (1999) The evolution of North American attitudes towards marine mammals. In: Twiss JR, Reeves RR (editors), Conservation and Management of

- Marine Mammals Washington, DC: Smithsonian Institution Press, pp 10–47
- Leighfield T (2002) Indian River Lagoon dolphin mortality. Unpublished report. 1 p
- Lipscomb TP, Kennedy S, Moffett D, Krafft A, Klaunberg A, Lichy JH, et al. (1996) Morbillivirus epizootic in bottlenose dolphins of the Gulf of Mexico. *Journal of Veterinary Diagnostic Investigation* 8:283–290
- Lipscomb TP, Schulman Y, Moffett D, Kennedy S (1994) Morbilliviral disease in Atlantic bottlenose dolphins (*Tursiops truncatus*) from 1987–1988 epizootic. *Journal of Wildlife Diseases* 30:567–571
- Loughlin TR (1994) Marine Mammals and the Exxon Valdez. San Diego, CA: Academic Press, 395 pp
- Mase B, Jones W, Ewing R, Bossart G, Van Dolah F, Leighfield T, et al. (2000) Epizootic in bottlenose dolphins in the Florida panhandle: 1999–2000. In: *Proceedings of the International Association for Aquatic Animal Medicine Annual Conference*, New Orleans, LA, 2000. p 593
- Mazet J, Torres de la Riva G, Gulland F, Langlois G, Kreuder C (2005) Marine mammal strandings associated with toxic algal blooms along the California coastline in 2002. Research Agreement Number DG133F-02-SE-0869
- McLellan WA, Friedlander AS, Mead JG, Potter CW, Pabst AP (2002) Analyzing 25 years of bottlenose dolphin (*Tursiops truncatus*) strandings along the Atlantic coast of the USA: do historic records support the coastal migratory stock hypothesis?. *Journal of Cetacean Research and Management* 4:297–304
- Miller MA, Gardner IA, Kreuder C, Paradies DM, Worcester KR, Jessup DA (2002) Coastal freshwater runoff is a risk factor for *Toxoplasma gondii* infection of southern sea otters (Enhydra lutris nereis).. International Journal of Parasitology 32:997–1006
- Moore SE, Urban JR, Perryman W, Gulland F, Peres-Cortes H, Wade P, et al. (2001) Are gray whales hitting "K" hard?. *Marine Mammal Science* 17:954–958
- National Research Council (1999) From Monsoons to Microbes. Understanding the Ocean's Role in Human Health. Washington, DC: National Academy Press, 132 pp
- O'Shea TJ, Rathburn GB, Bonde RK, Buergelt CD, Odell DK (1991) An epizootic of Florida manatees associated with a dinoflagellate bloom. *Marine Mammal Science* 7:165–179
- Osterhaus A, van de Bildt M, Vedder L, Martina B, Niesters H, Vos J, et al. (1998) Monk seal mortality: virus or toxin?. *Vaccine* 16:979–981
- Osterhaus ADME, Groen J, De Vries P, UytedHaag FGCM, Klingeborn B, Zarnke R (1988) Canine distemper virus in seals. *Nature* 335:403–404
- Osterhaus ADME, Groen J, UytdeHaag FGCM, Visser IKG, Van de Bildt MWG, Bergman A, et al. (1989) Distemper virus in Baikal seals. *Nature* 338:209–210
- Osterhaus ADME, Yang H, Spijkers HE, Groen J, Teppema JS, van Steenis G (1985) The isolation and partial characterization of a highly pathogenic herpesvirus from the harbor seal (*Phoca vitulina*).. Archives of Virology 86:239–251
- Preen AR, Marsh H (1995) Response of dugongs to large scale loss of sea grass from Hervey Bay, Queensland, Australia. *Wildlife Research* 22:507–519
- Raverty SA, Lambourn DM, Jeffries SJ, Cameron CE, Norman SA, Zuerner R, et al. (2005) Incursion of *Leptospira interrogans*, serovar *pomona* into the Pacific north west by migratory subadult and adult California sea lions (*Zalophus californianus*). In:

- Proceedings of the 36th Annual Conference of the International Association for Aquatic Animal Medicine, May 14–18, 2005, Seward, Alaska, pp 245–247
- Read A, Murray KY (2000) Gross evidence of human-induced mortality in small cetaceans. U.S. Department of Commerce, NOAA Technical Memorandum, NMFS-OPR-15, 21 pp
- Reidarson TH, McBain J, House C, King D, Stott JL, Krafft A, et al. (1998) Morbillivirus infection in stranded common dolphins from the Pacific ocean. *Journal of Wildlife Diseases* 34:771–776
- Ridgway SH (editor) (1996) Final Report from the Right Whale Necropsy Assessment Team: Results, Analysis, and Recommendations. NRaD TD 2934, 51 pp
- Ross HM, Jahans KL, MacMillan AP, Reid RJ, Thompson PM, Foster G (1996) Brucella species infection in North Sea seal and cetacean populations. *Veterinary Record* 138:647–648
- Ross PS, DeSwart RL, Reijnders PJH, Van Loveren H, Vos JG, Osterhaus ADME (1995) Contaminant-related suppression of delayed-type hypersensitivity and antibody responses in harbor seals fed herring from the Baltic sea. *Environmental Health Perspectives* 103:162–167
- Scholin CA, Gulland F, Doucette GJ, Benson S, Busman M, Chavez FP, et al. (2000) Mortality of sea lions along the central California coast linked to a toxic diatom bloom. *Nature* 403:80–84
- Schulman Y, Lipscomb TP, Moffett D, Krafft AE, Lichy JH, Tsai MM, et al. (1997) Histologic, immunohistochemical, and polymerase chain reaction studies of bottlenose dolphins from the 1987–1988 United States Atlantic coast epizootic. *Veterinary Pathology* 34:288–295
- Scott GP, Burn DM, Hansen LJ (1988) The dolphin die-off: long-term effects and recovery of the population. In: *Proceedings of the Oceans '88 Conference*, Baltimore, MD, pp 819–823
- Sergeant DE (1982) Mass strandings of toothed whales (Odontoceti) as a population phenomenon. Scientific Report of the Whale Research Institute 34:1

- Silvagni PA, Lowenstine LJ, Spraker T, Lipscomb TP, Gulland FM (2005) Pathology of domoic acid toxicity in California sea lions (*Zalophus californianus*). *Veterinary Pathology* 42:184–191
- Stoddard R, Gulland FMD, Atwill ER, Lawrence J, Jang S, Conrad PA (2005) Salmonella and Campylobacter spp. in northern elephant seals, California. Emerging Infectious Diseases 11:1967– 1969
- Sweeney J (1992) Veterinary assessment report, Tursiops truncatus, Matagorda Bay, Texas, July 1992. Contract Report, NOAA-NMFS, SEFSC, Contribution MIA-92/93-41. 10 pp. + appendices
- Thompson D, Lonergan ME, Duck C (2005) Population dynamics of harbour seals (*Phoca vitulina*) in England: growth and catastrophic declines. *Journal of Applied Ecology* 42:638–648
- Thompson PM, Hall AJ (1993) Seals and epizootics—what factors might affect the severity of mass mortalities?. *Mammal Review* 23:149–154
- Trillmich F, Ono KA (1991) *Pinnipeds and El Niño*. New York: Springer-Verlag, 293 pp
- Van Dolah FM (2000) Marine algal toxins: origins, health effects, and their increased occurrence. *Environmental Health Perspectives* 108(Suppl 1):133–141
- Vos DJ, Sheldon KEW (2005) Unusual mortality in the depleted Cook Inlet Beluga (*Delphinapterus leucas*) population. *Northwestern Naturalist* 86:59–65
- Ward JR, Lafferty KD (2004) The elusive baseline of marine diseases: are diseases in ocean ecosystems increasing?. PLoS Biology 2:0542–0547
- Wilkinson DM (1996) National contingeny plan for response to unusual marine mammal mortality events. NOAA Technical Memorandum NMFSOPR-9, September 1696, Silver Spring, MD